

OPTIMIZING CHIRALITY AND SCALING OF LATTICE FERMIONS

W. BIETENHOLZ
 NORDITA, Blegdamsvej 17
 DK-2100 København Ø, Denmark
 NORDITA-2000-1-HE, hep-lat/0001001

1. Introduction

If we construct a lattice fermion formulation, there are a number of goals to be considered: *doubling* should be avoided; even at finite lattice spacing a , we want to represent *chiral symmetry* in a sound way; and we are seeking a good *scaling* behavior. Conceptually we have to require *locality* (the lattice Dirac operator $D(x, y, U)$ has to decay at least exponentially in $|x - y|$). In addition, for practical purposes we desire a high level of locality, i.e. a *fast* exponential decay or even *ultralocality* (which means that the couplings in D drop to zero beyond a finite number of lattice spacings). A further issue is a good approximation to rotation invariance. Last but not least, the formulation should be simple enough to allow for efficient simulations. Here we report on a construction, which is designed to do justice to all of these goals.

2. Ginsparg-Wilson fermions (an unconventional introduction)

For a lattice Dirac operator D , full chiral invariance ($\{D, \gamma_5\} = 0$) is incompatible with other basic requirements (Hermiticity, locality, absence of doublers, discrete translation invariance) [1]. Therefore we only implement a *modified chiral symmetry*, which does allow D to fulfill those requirements. For such a modified chiral transformation we start from the ansatz

$$\bar{\psi} \rightarrow \bar{\psi} (1 + \epsilon[1 - F]\gamma_5) , \quad \psi \rightarrow (1 + \epsilon\gamma_5[1 - G]) \psi , \quad (1)$$

ϵ being an infinitesimal transformation parameter. The transformation, and therefore F and G should be local, and $F, G = O(a)$, so that we reproduce the full chiral symmetry in the (naive) continuum limit.¹ Invariance of the

¹For convenience, we set $a = 1$ in the formulae (on an isotropic Euclidean lattice), but we classify the terms nevertheless by the order of a that they (would) belong to.

Lagrangian $\bar{\psi}D\psi$ holds to $O(\epsilon)$ if ²

$$\{D, \gamma_5\} = F\gamma_5 D + D\gamma_5 G. \quad (2)$$

This implies a continuous modified chiral symmetry, which has the full number of generators. It may be compared to the remnant chiral symmetry of staggered fermions: there the doubling problem is not solved, and one is only left with a $U(1) \otimes U(1)$ symmetry, which does, however, protect the mass from additive renormalization. The same can be shown here if we assume “ γ_5 -Hermiticity”, $D^\dagger = \gamma_5 D \gamma_5$, and we choose $F = DR$, $G = RD$, where R is *local* again, non-trivial and $[R, \gamma_5] = 0$ (this generalizes Ref. [2]). Then eq. (2) turns into the *Ginsparg-Wilson relation* (GWR) [3] ³

$$D + D^\dagger = 2D^\dagger R D, \quad (3)$$

and it implies the absence of additive mass renormalization (see also Ref. [5]) since

$$(\sqrt{2RD}^\dagger \sqrt{2R} - 1)(\sqrt{2RD} \sqrt{2R} - 1) = 1. \quad (4)$$

As another illustration we can write the GWR as $\{D^{-1}, \gamma_5\} = 2R\gamma_5$, and we see that a local term R does not shift the poles in D^{-1} (in contrast to the cases where $\{D, \gamma_5\}/2$ is local, such as a mass or a Wilson term). ⁴ [With respect to the general ansatz, we have to require the right-hand side of $D^{-1} + D^{\dagger -1} = D^{\dagger -1} \gamma_5 F \gamma_5 + G D^{-1} = D^{-1} F + \gamma_5 G \gamma_5 D^{\dagger -1}$ to be local.]

γ_5 -Hermiticity implies $R^\dagger = R$. If we now start from some lattice Dirac operator D_0 (obeying the assumption of the Nielsen-Ninomiya theorem such as absence of doublers, but otherwise quite arbitrary), we can construct a Ginsparg-Wilson operator D from it by enforcing eq. (4) as

$$D = \frac{1}{\sqrt{2R}} \left[1 + \frac{A}{\sqrt{A^\dagger A}} \right] \frac{1}{\sqrt{2R}}, \quad A := \sqrt{2R} D_0 \sqrt{2R} - 1. \quad (5)$$

This is the generalization [7] of the “*overlap formula*”, which uses the “*standard GW kernel*”

$$R_{x,y}^{(st)} := \frac{1}{2} \delta_{x,y}, \quad (6)$$

²In our short-hand notation, the ‘products’ are convolutions in c-space.

³ γ_5 -Hermiticity is essentially inevitable for any sensible solution, but if we want to formulate the GWR even without this assumption, then it reads $\{D, \gamma_5\} = 2DR\gamma_5 D$. This follows from the immediately obvious prescription $\{D^{-1}, \gamma_5\} = 2R\gamma_5$. Alternatively, if we require D to be normal we arrive at $\{D, \gamma_5\} = 2RD\gamma_5 D$ [4]. However, for the results presented in Sec. 3, 4 this doesn’t matter, since we always use $R_{x,y} \propto \delta_{x,y}$.

⁴An analogous treatment is also conceivable in the continuum. For comments related to the Pauli-Villars regularization, see Ref. [6]. In dimensional regularization γ_5 is a notorious trouble-maker. It may be useful to substitute it by operators $(1 - F)\gamma_5$ resp. $\gamma_5(1 - G)$ at the suitable places. In $d + \varepsilon$ dimensions (d even), F and G could take the form $\varepsilon DR/\mu$ resp. $\varepsilon RD/\mu$ (where R is some local term, and μ is the usual scale in dimensional regularization). We expect the chiral anomaly to be reproduced correctly as $\varepsilon \rightarrow 0$.

and which leads from the Wilson fermion $D_0 = D_W$ to the Neuberger fermion $D = D_{Ne}$ [8]. In the general solution of the GWR, eq. (5), we can obviously vary the parameters in D_0 or in R (or in both) in many ways, without violating their required properties. This shows that there exists a continuous set of GWR solutions D in the space of coupling parameters.

Any solution of the GWR is related to a fully chirally invariant Dirac operator $D_\chi = D(1 - RD)^{-1}$, which is, however, non-local (in the free case, $D_\chi(p)$ has poles, cf. eq. (4)). Vice versa, if we start from some D_χ with this type of non-locality (such as the Rebbi fermion [9], for example) we can construct a GW solution [7, 10] $D = D_\chi(1 + RD_\chi)^{-1}$, which is local, at least in the free and weakly interacting case. The mechanism of providing locality by inserting a local term $R \neq 0$ is known from the framework of perfect actions, where the factor R^{-1} occurs in a Gaussian block variable renormalization group transformation term of the fermions [3, 11]. Hence $R \rightarrow 0$ corresponds to a δ function block variable transformation, and the corresponding perfect action has a Rebbi-type non-locality. The transition to locality requires the superficial breaking of the full chiral symmetry, $R \neq 0$: chirality is manifest in the action only in the sense of the GWR, but it is fully present in the physical observables [12, 13].⁵ In contrast to the Rebbi fermion [16], the axial anomaly is correctly reproduced [17] for the perfect action at any local term R , including the perfect D_χ (for $R = 0$). This should also be checked if one generally wants to use D_χ in an indirect way [18], by measuring the right-hand side of $\langle D_\chi^{-1} \rangle = \langle D^{-1} \rangle - \langle R \rangle$.

By introducing a non-trivial kernel R we have relaxed the condition of chiral symmetry somewhat — without doing harm to the physical properties related to chirality [3, 5, 2] — and this allows for locality of D (as well as the absence of doublers etc.)⁶ without contradiction to the Nielsen-Ninomiya theorem. In the case of the Neuberger fermion D_{Ne} , locality has been demonstrated in a smooth gauge background. In particular, zero eigenvalues in $A^\dagger A$ are excluded if the inequality (in d dimensions)

$$\|1 - P\| < \frac{b}{d(d-1)} \quad (7)$$

holds for any plaquette variable P and a suitable bound b . From Ref. [19] we obtain $b = 0.4$, which has recently been improved to $b = (1 + 1/\sqrt{2})^{-1} \simeq 0.586$ [20]. Still, this constraint is somewhat inconvenient; for instance, at least one eigenvalue of $A^\dagger A$ has to cross zero if we want to change the topological sector.

⁵Such a superficial symmetry breaking in the transformation term is *not* necessary in order to preserve *supersymmetry* in a RGT, and hence in a perfect action [14]. A. Thimm also commented on an analogous treatment of further symmetries [15].

⁶This holds at least as long as the gauge background is smooth. At very strong coupling, locality is uncertain, and also the doubling problem can return, see Subsec. 4.2.

Furthermore, the GWR allows for locality only in the sense that the couplings in $D(x, y, U)$ decay exponentially, but *not* for ultralocality. This was conjectured intuitively in Ref. [7]. To demonstrate this No-Go rule for GW fermions, it is sufficient to show it for the free fermion. A proof, which was specifically restricted to $R = R^{(st)}$, has been given in Ref. [21]. By now, a complete proof covering *all* (local) GW kernels R has been added, hence this rule is completely general [22].

In that context, it is amusing to reconsider the ordinary Wilson fermion. From the mass shift we know that it is certainly not a GW fermion in general, and according to our No-Go rule not even the free Wilson fermions can obey any GWR. Indeed, if we insert the free D_W into the GWR and solve for $R_{x,y}$, we find that it decays $\propto |x - y|^{-4}$ in $d = 2$, and $\propto |x - y|^{-6}$ in $d = 4$, which is nonlocal and therefore not a GW kernel.

The exponential decay of D is satisfactory from the conceptual point of view, but the existence of couplings over an infinite range is a problem for practical purposes. One would hope for a *high degree of locality* at least, i.e. for a fast exponential decay. Almost all the literature on overlap fermions solely deals with the Neuberger fermion, but it turns out that the couplings in D_{Ne} do not decay as fast as one would wish, see Fig. 5. Moreover, also other properties listed in Sec. 1 — most importantly scaling, but also approximate rotational invariance — are unfortunately rather poor. This is obvious even from the free fermion, see Figs. 2, 5. On the level of the action, there are generally no $O(a)$ artifacts for GW fermions, because any additional clover term violates the GWR [23], but from the free case we see already that the $O(a^2)$ scaling artifacts in D_{Ne} are large.⁷ And yet its simulation is tedious (the quenched case requires already a similar effort as simulating D_W with dynamical fermions [25]), allowing only for the use of small lattices.

However, D_{Ne} arises only from a very special choice in a large class of GWR solutions described by eq. (5), namely $D_0 = D_W$ and $R = R^{(st)}$. The message of this report is that there are better options, and we are going to show in Sec. 3, 4 how *improved overlap fermions* can be constructed, tested and applied. This generalization and the improvement concept for overlap fermions was introduced in Ref. [7]. It was extensively tested in the framework of the Schwinger model [26].

⁷Our study of the 2d and 4d free fermion, as well as the 2d interaction fermion, show consistently that the artifacts in D_{Ne} are even worse than those in D_W . Ref. [24] is more optimistic about the scaling of the Neuberger fermion (but it does not include a comparison to other types of lattice fermions).

2.1. THE CONCEPT OF IMPROVING OVERLAP FERMIONS

We first summarize the main idea: if D_0 happens to be a GW operator already (with respect to a fixed term R), then eq. (5) yields $D = D_0$; the operator reproduces itself. Therefore, any GW fermion — such as the perfect [3] or the classically perfect [5] fermion — is automatically an overlap fermion too. The construction of a classically perfect action for asymptotically free models only requires minimization — no (numeric) functional integral, in contrast to the perfect action — and it still has the additional virtues of excellent scaling and rotation invariance. Unfortunately, a really powerful quasi-perfect action is not available so far for interacting fermions in $d = 4$.

However, if we manage to construct at least an *approximate GW fermion*, then we can expect it to *change only modestly* if we insert it as D_0 in the overlap formula (for the corresponding R), $D \approx D_0$. If our approximate GW fermion is in addition short-ranged, then we can expect D to have a high degree of locality, since the long distance couplings are turned on just a little in D . (Also $D_0 = D_W$ is short-ranged, but since this is far from a GW fermion, it changes a lot in the overlap formula, and those long distance couplings cannot be predicted to be tiny). Similarly, if D_0 scales well, then we can expect this quality to be essentially preserved in D if $D \approx D_0$, and the same argument applies to the approximate rotation invariance. In Sec. 3 we discuss examples for promising approximate GW operators. In Sec. 4 they are transformed into exact GW fermions, and the above predictions are verified.

3. Short-ranged approximate Ginsparg-Wilson fermions

Perfect free fermions can be constructed and parameterized in c-space explicitly [13]. If we choose the term R such that the locality is optimal at mass zero, then we arrive at the standard form $R^{(st)}$. To make such a fermion tractable, first of all its couplings have to be truncated to a short range. A truncation to couplings inside a unit hypercube (“*hypercube fermion*”, HF) was performed in Ref. [27] by means of periodic boundary conditions. (Alternative 4d truncated perfect HFs can be found in Refs. [28].) Truncation causes some scaling artifacts, but they are small, so that the free HF is still strongly improved over the Wilson fermion. This can be observed from the dispersion relation as well as thermodynamic scaling ratios [27, 29]. At the same time, truncation also implies a (small) violation of the GWR, in agreement with the absence of ultralocal GW fermions. The spectrum of a GW fermion with $R_{x,y} = \delta_{x,y}/(2\mu)$, ($\mu > 0$) is situated on a circle in $\mathcal{C}$ with center and radius μ (GW circle), see eq. (4). Hence we can test the quality of our approximate GW fermion (with respect to $R^{(st)}$) by checking how

close its spectrum comes to a unit circle. This is shown in Fig. 1 for the 4d HF of Ref. [27] on a 20^4 lattice. We see that we have a good approximation, especially in the physically important regime of eigenvalues close to 0.




Figure 1. The spectrum of a truncated perfect, free HF on a 20^4 lattice (plotted in $\mathbb{C}$).

In $d = 2$ we start off from a similar massless HF, which is optimized for its scaling behavior; its set of couplings is given in Ref. [26], and its strong improvement over the Wilson fermion is visible in Fig. 2.⁸ The free spectrum of the “*scaling optimal hypercube fermion*” (SO-HF) is again close to a unit circle, on the same level as the 4d spectrum in Fig. 1. There are also other ways to see that we are in the vicinity of a GW fermion, for instance by summing over its violation (squared) in each site [26], or by inserting the HF into the GWR and solving for R : for the SO-HF, $R_{x,y}$ decays about 6 times faster than the corresponding (pseudo-) R for D_W . If we compare the truncated perfect HF to D_W , then this factor even amounts to ≈ 71 in $d = 2$, and to ≈ 75 in $d = 4$.

We now proceed to the 2-flavor Schwinger model, and we gauge the SO-HF by attaching the couplings to the shortest lattice paths only. When there are several shortest paths, the coupling is split and attached to them in equal parts. Moreover we add a clover term with coefficient 1. For the

⁸As an alternative, one could directly optimized the chiral properties by minimizing the violation of the GWR [7, 26, 30]. However, those properties can still be corrected later on by means of the overlap formula, whereas no such tool is available to correct the scaling behavior. Hence it is important to optimize *scaling* — rather than *chirality* — from the beginning.

Moreover, if one constructs a truncated approximate GW fermions in a short range, there is no direct control over the level of locality of the full GW fermion that one approximates. Actually it is not even certain if the GW solution that one would obtain by gradually extending the range to infinity is local at all, since also non-local solutions of the GWR exist. A simple example for that can be constructed from the SLAC fermion [31]. The Dirac operator $D = D_{SLAC}(1 + RD_{SLAC})^{-1}$ solves the GWR, but it is non-local.




Figure 2. Scaling of free fermions: the fermion dispersion relation (left) and the thermodynamic scaling ratio $\text{pressure}/(\text{chemical potential})^2$ at zero temperature (right).

gauge part we use the standard plaquette action (which is perfect for 2d pure $U(1)$ gauge theory [13]).

This simple “gauging by hand” causes a further deviation from the GWR, which is increasingly manifest if the gauge background becomes rougher. In Fig. 3 we show the spectra for typical configurations at $\beta = 2$ and at $\beta = 6$. It turns out, however, that the SO-HF is indeed an approx-




Figure 3. The spectra of the 2d scaling optimal hypercube fermion (SO-HF) for typical configurations at strong resp. weak coupling, approximating a GW circle.

imate GW fermion up to a considerable couplings strength. Next we test the scaling behavior in the presence of gauge interaction. Our simulation results (here and below) were obtained in collaboration with I. Hip. They are based on 5000 quenched configurations on a 16×16 lattice at $\beta = 6$ (we use the same set of configurations for all types of fermions), but the evaluation does include the fermion determinant, following Ref. [32]. Fig. 4 shows the dispersion relations for the two meson-type states, a massless triplet and one massive mode [33], which we denote as π and η (by analogy). Again the SO-HF is drastically improved over the Wilson fermion (at $\kappa_c = 0.25927$, from Ref. [34]). It even reaches the same level as a (very

mildly truncated) classically perfect action [35], which was parameterized by 123 independent couplings per site, whereas only 6 such couplings are used for the SO-HF. Therefore, it is realistic to extend the HF formulation to QCD, and in fact it has been shown already that minimally gauged 4d HFs can indeed be applied in QCD simulations [36, 37]. However, in QCD as well as in the Schwinger model, we observed as an unpleasant feature of the directly applied HF a strong additive mass renormalization. Using the SO-HF, even at $\beta = 6$ the π mass is renormalized from 0 to 0.13, as Fig. 4 shows. This corresponds to a lowest real eigenvalue around 0.03, cf. Fig. 3. As an even more striking example, the 4d massless HF, minimally gauged by hand and applied to QCD at $\beta = 5$, leads to a “pion mass” of 3.0 [27], and at $\beta = 6$ the critical bare HF mass amounts to -0.92 [37] (which is unfavorable for the level of locality before truncation, and hence for the magnitude of the truncation effects). We overcome this problem in the next section by inserting the SO-HF into the overlap formula.

[As a further alternative to negative bare mass and overlap, we can reach the exact chiral limit solely by the use of fat links [26]. This is essentially equivalent to gauging the HF such that the interacting GWR is just violated modestly. As yet another possibility one may attach an amplification factor $\gtrsim 1$ to each link [38].]




Figure 4. The “meson” dispersion relations for some types of 2d fermions in the 2-flavor Schwinger model at $\beta = 6$.

4. Improved overlap fermions

We now perform the second step in our program and insert the SO-HF — which is an approximate GW fermion with an excellent scaling behavior — into the overlap formula (5) with $R = R^{(st)}$ given in eq. (6). This leads to an exact GW fermion and therefore to all the nice properties related to chirality, which are extensively discussed in the recent literature on GW fermions: correct anomalies; no renormalization of mass zero, vector current and flavor non-singlet axial vector current; no mixing of weak ma-

trix elements; no exceptional configurations [5]; correctly reproduced chiral symmetry breaking [39] etc.

Our first prediction was that the level of locality should be improved over the Neuberger fermion, and this is clearly confirmed, both, in the free and in the interacting case. Fig. 5 compares the decay of the couplings of the free fermion (left) and the decay of the “maximal correlation” f over a certain distance r — as suggested in Ref. [19] — at $\beta = 6$ (right).⁹ (In the latter figure one could still try to improve the locality in both cases by deviating from $R^{(st)}$ and tuning R for optimal locality.) Next we want




Figure 5. The level of locality for the Neuberger fermions vs. overlap SO-HF: the decay of the free couplings in all directions in infinite volume (left) and the decay of the “maximal correlation” over a distance r , as defined in Ref. [19], on a 24×24 lattice at $\beta = 6$ (right). The figure on the left also illustrates how well rotation invariance is approximated.

to verify if the good scaling quality survives the modification due to the overlap formula. For the free fermions, this is confirmed in an impressive way, see Fig. 2.¹⁰ On fine lattices, the scaling is practically identical before and after the use of the overlap formula; only as the lattice becomes really coarse, the overlap does some harm to the scaling quality at some point.

In the presence of gauge interaction, we consider again the “meson” dispersions, and we observe again the persistence of the improvement, see Fig. 6.

Furthermore, continuous rotation invariance is approximated much better for the HF than for the Wilson fermion (after all, also this holds *exactly* for the perfect fermion [13]). In Ref. [26] this was also tested in the interacting case by measuring how smoothly correlations decay with the Euclidean distance, and we observed that the SO-HF is again by far superior over

⁹One puts a unit source at some site x and defines $f(r) := \max_y \{ \|\psi(y)\| \mid |x-y| = r \}$.

¹⁰The dispersion curves of the overlap fermions just stop inside the Brillouin zone due to the square root. The end-points can be shifted by choosing $R_{x,y} = \delta_{x,y}/(2\mu)$ and varying μ . We show $\mu = 1$ which is a reasonable choice; if we push the end-point to far, the danger of doubling increases. On the other hand, the regime of small momenta is needed, of course.




Figure 6. The “meson” dispersion relations for two types of overlap fermions at $\beta = 6$.

the Wilson fermion; once more it reaches the same level as the classically perfect action. The overlap formula (5) suggests that this property is essentially inherited for the overlap fermions, and indeed we observed a similar level for the Wilson fermion and the Neuberger fermion on one hand, and for the SO-HF and the overlap SO-HF on the other hand. This confirms that also the strongly improved approximate rotational invariance of the SO-HF survives if it is turned into an overlap fermion. For the free fermion, this progress can also be observed in Fig. 5 (left) from the width of the “cones”.

4.1. CHIRAL CORRECTION IN TERMS OF A POWER SERIES

In $d = 2$ we can afford an exact evaluation of the notorious square root in the overlap formula, but in QCD this is not feasible any more. In the recent literature, a number of iterative procedures have been suggested for the Neuberger fermion [19, 40]. In Ref. [26] we presented a new method, which is very simple and robust, and which is especially designed for the case where D_0 is an approximate GW fermion already. We evaluate the square root in

$$D = 1 + \frac{A}{\sqrt{A^\dagger A}}, \quad A := D_0 - 1 \quad (8)$$

as a power series in $\varepsilon := A^\dagger A - 1$. For $D_0 = D_{HF}$ we have $\|\varepsilon\| \ll 1$ (if the configuration is not extremely rough), hence the expansion converges rapidly. On the other hand, for the case of the Neuberger fermion, i.e. for $D_0 = D_W$, this expansion fails to converge even in the free case, which is presumably the reason why it had not been considered in the earlier literature. We call this method a “perturbative chiral correction”, where the perturbative expansion refers to the GWR violation ε (and not to the coupling g). It yields a Dirac operator of the form

$$D_{p\chi c} = 1 + AY. \quad (9)$$

For the chiral correction to $O(\varepsilon^n)$, Y is a polynomial in $A^\dagger A$ of order n , for instance,

$$\begin{aligned} Y &= [3 - A^\dagger A]/2 & \text{for } n = 1, \\ Y &= [15 - 10A^\dagger A + 3(A^\dagger A)^2]/8 & \text{for } n = 2, \text{ etc.} \end{aligned} \quad (10)$$

Hence the computational effort amounts roughly to $1 + 2n$ matrix-vector multiplication (the matrix being A or $A^\dagger$), i.e. it increases *only* linearly (though the convergence is also just linear). Actually this represents a fermion with couplings of range $1 + 2n$ in each component, and it would be very tedious to implement it explicitly, even for $n = 1$. However, due to its specific form we never need to do so; D_{pxc} can always be evaluated by iteration of the above matrix-vector products.

The crucial question now is if the first few orders are sufficient already to do most of the chiral correction. We first look at the free SO-HF, and Fig. 7 (left) shows that the first order alone does practically the full job. In this context, we also obtain a geometric picture of the effect of the overlap formula: for $R_{x,y} = \delta_{x,y}/(2\mu)$ it can be viewed as a *projection* of the eigenvalues onto the circle with center and radius μ . This projection is often close to radial.




Figure 7. The convergence of the SO-HF to a GW fermion under perturbative chiral correction (on a 16×16 lattice): the free fermion spectrum (left) and the distribution of the small real eigenvalues at $\beta = 6$, moving rapidly closer to 0 (right).

In the interacting case, the convergence to the circle under iteration is slowest in the arc around 0. As an example, we show in Fig. 7 (right) a histogram of the small real eigenvalues at $\beta = 6$. We see that the mass renormalization is removed almost completely if we proceed to $O(\varepsilon^2)$.

4.2. BEHAVIOR IN EXTREMELY ROUGH GAUGE CONFIGURATIONS

For smooth configurations, it is easy to find a parameter μ — and hence a GW circle where the spectrum is mapped on — such that the small (large)

real eigenvalues are mapped on 0 (2μ). This leaves the index unchanged (though it is defined by exact zero modes now) and it provides a sensible definition of the topological charge via the index theorem [5, 2]. It also means that the doubling problem is safely avoided for all typical configurations at moderate or large β .

However, if we dare proceeding to extremely strong coupling, then it is not possible any more to find such a center μ of a GW circle, which does the right mapping for all typical configurations. In fact, for extremely rough QCD test configurations (on very small lattices) it was observed explicitly that *all* the eigenvalues of the minimally gauged D_{HF} are close to the arc of the GW circle, which is opposite to 0 [41]. For D_W the eigenvalues are densely scattered over a wide area with a large real part. Examples for such spectra of an extremely rough configuration are shown in Fig. 8, which was provided by N. Eicker, I. Hip and Th. Lippert. At a coupling strength where such configurations are frequent, the doubling problem is back for those overlap fermions, which are constructed from some simple D_0 (for the Schwinger model, this problem sets in around $\beta \approx 1$ [41]). This agrees with the result of a strong coupling expansion (in the Hamiltonian formulation) which applies to D_{Ne} [42].¹¹ In such cases, the construction of D_{Ne} maps all (almost) real eigenvalues onto the arc close to 2, hence additive mass renormalization is back as well — again in agreement with Ref. [42]. In view of the latter point one could be tempted to just use a large mass parameter $\mu > 1$; this does not help, however, with respect to the doubling problem. At very strong coupling, only the (classically) perfect action would help.

As we mentioned earlier, also locality is in danger in that regime [19], and we should therefore keep away from it. As one more advantage of choosing D_0 to be an approximate GW fermion, the regime of β where we are (statistically) on safe grounds is enlarged compared to the Neuberger fermion.

Of course, in the safe regime where β is large enough (in QCD this includes $\beta = 6$ for sure [19]) the chiral correction of $D_0 = D_{HF}$ can also be carried out by iteration methods different from the one described in the previous Subsection, for alternative experiments see Ref. [44]. The efficiency in QCD is still to be compared, but for sure in any method the convergence will be much faster for $D_0 = D_{HF}$ than for $D_0 = D_W$.

5. Conclusions

Our program outlined in Sec. 2.1 has been realized in the Schwinger model, and the properties of a resulting improved overlap fermion have been tested

¹¹For a Euclidean strong coupling hopping parameter expansion of D_{Ne} , see Ref. [43].




Figure 8. The fermion spectrum for an extremely rough QCD test configuration on a 4^4 lattice: for the hypercube fermion of Ref. [27] (left), and for the Wilson fermion (right).

extensively. They are all clearly superior over the Neuberger fermion, confirming our prediction: the overlap SO-HF scales much better, it is more local and it comes much closer to rotation invariance.

The question now is the applicability of this program in $d = 4$. The 4d HF formulation is worked out already, and the corresponding improved overlap fermion is currently under investigation in QCD by the SESAM collaboration in Jülich and Wuppertal.

Acknowledgment Most of the results presented here are based on my collaboration with Ivan Hip, and I would like to thank him for his crucial contributions. Furthermore I am indebted to P. Damgaard, T. DeGrand, P. Hernández, A. Hoferichter, K. Jansen, J. Jersák, F. Klinkhamer, Th. Lippert, K.F. Liu, M. Lüscher, J. Nishimura, P. Rakow, K. Schilling and A. Thimm for useful comments. Finally I would like to thank the organizers of this work-shop, in particular V. Mitrjushkin, for their kind hospitality in beautiful Dubna.

References

1. H. Nielsen and M. Ninomiya, *Nucl. Phys.* **B185** (1981) 20.
2. M. Lüscher, *Phys. Lett.* **B428** (1998) 342.
3. P. Ginsparg and K. Wilson, *Phys. Rev.* **D25** (1982) 2649.
4. W. Kerler, **hep-lat/9905010**.
5. P. Hasenfratz, V. Laliena and F. Niedermayer, *Phys. Lett.* **B427** (1998) 125.
P. Hasenfratz, *Nucl. Phys.* **B525** (1998) 401.
6. K. Fujikawa, *Nucl. Phys.* **B546** (1999) 480.
7. W. Bietenholz, *Eur. Phys. J.* **C6** (1999) 537.
8. H. Neuberger, *Phys. Lett.* **B417** (1998) 141; **B427** (1998) 353.

9. C. Rebbi, *Phys. Lett.* **B186** (1987) 200.
10. T.-W. Chiu and S. Zenkin, *Phys. Rev.* **D59** (1999) 074501.
11. U.-J. Wiese, *Phys. Lett.* **B315** (1993) 417.
W. Bietenholz and U.-J. Wiese, *Nucl. Phys. B (Proc. Suppl.)* **34** (1994) 516.
12. W. Bietenholz and U.-J. Wiese, *Nucl. Phys. B (Proc. Suppl.)* **47** (1996) 575.
13. W. Bietenholz and U.-J. Wiese, *Nucl. Phys.* **B464** (1996) 319.
14. W. Bietenholz, *Mod. Phys. Lett.* **A14** (1999) 51.
15. A. Thimm, talks presented at LATTICE'99 and at this workshop.
16. G. Bodwin and E. Kovacs, *Phys. Lett.* **B193** (1987) 283; *Phys. Rev.* **D35** (1987) 3198. M. Campostrini, G. Curci and A. Pelissetto, *Phys. Lett.* **B193** (1987) 273.
A. Pelissetto, *Ann. Phys.* **182** (1988) 177.
17. W. Bietenholz and U.-J. Wiese, *Phys. Lett.* **B378** (1996) 222.
18. S. Capitani et al., *Phys. Lett.* **B468** (1999) 150.
19. P. Hernández, K. Jansen and M. Lüscher, *Nucl. Phys.* **B552** (1999) 363.
20. H. Neuberger, **hep-lat/9911004**.
21. I. Horváth, *Phys. Rev. Lett.* **81** (1998) 4063.
22. W. Bietenholz, **hep-lat/9901005**.
23. F. Niedermayer, *Nucl. Phys. (Proc. Suppl.)* **73** (1999) 105.
24. K.F. Liu, S.J. Dong, F.X. Lee and J.B. Zhang, **hep-lat/9909061**.
25. P. Hernández, K. Jansen and L. Lellouch, **hep-lat/0001008**.
26. W. Bietenholz and I. Hip, **hep-lat/9902019**, to appear in *Nucl. Phys. B*; **hep-lat/9908012**.
27. W. Bietenholz, R. Brower, S. Chandrasekharan and U.-J. Wiese, *Nucl. Phys. (Proc. Suppl.)* **53** (1997) 921.
28. F. Niedermayer, *Nucl. Phys. (Proc. Suppl.)* **53** (1997) 56.
W. Bietenholz, **hep-lat/9911015**, to appear in *Int. J. Mod. Phys. A*.
29. W. Bietenholz and U.-J. Wiese, *Phys. Lett.* **B426** (1998) 114.
30. C. Gattringer and I. Hip, **hep-lat/0002002**.
31. S. Drell, M. Weinstein and S. Yankielowicz, *Phys. Rev.* **D14** (1976) 487; 1627.
32. F. Farchioni, I. Hip and C.B. Lang, *Phys. Lett.* **B443** (1998) 214.
33. I. Sachs and A. Wipf, *Helv. Phys. Acta* **65** (1992) 652.
34. I. Hip, C.B. Lang and R. Teppner, *Nucl. Phys. B (Proc. Suppl.)* **63** (1998) 682.
35. C.B. Lang and T. Pany, *Nucl. Phys.* **B513** (1998) 645.
36. K. Orginos et al., *Nucl. Phys. (Proc. Suppl.)* **63** (1998) 904.
T. DeGrand, *Phys. Rev.* **D58** (1998) 094503.
37. N. Eicker et al., *Comput. Phys. Commun.* **119** (1999) 1.
38. N. Eicker et al., *in preparation*.
39. S. Chandrasekharan, *Phys. Rev.* **D60** (1999) 074503.
Y. Kikukawa and A. Yamada, *Nucl. Phys.* **B547** (1999) 413.
40. H. Neuberger, *Phys. Rev. Lett.* **81** (1998) 4060.
A. Boriçi, *Phys. Lett.* **B453** (1999) 46.
C. Liu, *Nucl. Phys.* **B554** (1999) 313.
41. I. Hip, *private communication*.
42. R. Brower and B. Svetitsky, **hep-lat/9912019**.
43. I. Ichinose and K. Nagao, **hep-lat/9910031**.
44. T. DeGrand, **hep-lat/9908037**.